

VOLUME LXXVI, ISSUE 5 PART I, MAY 2023

ISSN 0043-5147

E-ISSN 2719-342X

Wiadomości Lekarskie Medical Advances



Official journal of Polish Medical Association has been published since 1928



INDEXED IN PUBMED/MEDLINE, SCOPUS, EMBASE, EBSCO, INDEX COPERNICUS,
POLISH MINISTRY OF EDUCATION AND SCIENCE, POLISH MEDICAL BIBLIOGRAPHY

VOLUME LXXVI, ISSUE 5 PART I, MAY 2023

ISSN 0043-5147
E-ISSN 2719-342X

Wiadomości Lekarskie Medical Advances



Official journal of Polish Medical Association has been published since 1928



ALUNA Publishing House



Memory of
dr Władysław
Biegański

Wiadomości Lekarskie is abstracted and indexed in: PUBMED/MEDLINE, SCOPUS, EMBASE, INDEX COPERNICUS,
POLISH MINISTRY OF EDUCATION AND SCIENCE, POLISH MEDICAL BIBLIOGRAPHY

Copyright: © ALUNA Publishing House.

Articles published on-line and available in open access are published under Creative Common Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

The journal Wiadomości Lekarskie is cofinanced under Contract No.RCN/SN/0714/2021/1
by the funds of the Minister of Education and Science



Wiadomości Lekarskie Medical Advances

Editor in-Chief:

Prof. Władysław Pierzchała

Deputy Editor in-Chief:

Prof. Aleksander Sieroń

Statistical Editor:

Dr Lesia Rudenko

Managing Editor:

Agnieszka Rosa – amarosa@wp.pl

International Editorial Office:

Nina Radchenko (editor)

– n.radchenko@wydawnictwo-aluna.pl

Polish Medical Association (Polskie Towarzystwo Lekarskie):

Prof. Waldemar Kostewicz – President PTL

Prof. Jerzy Woy-Wojciechowski – Honorary President PTL

International Editorial Board – in-Chief:

Marek Rudnicki

Chicago, USA

International Editorial Board – Members:

Kris Bankiewicz	San Francisco, USA	George Krol	New York, USA
Christopher Bara	Hannover, Germany	Krzysztof Łabuzek	Katowice, Poland
Krzysztof Bielecki	Warsaw, Poland	Jerzy Robert Ładny	Białystok, Poland
Zana Bumbuliene	Vilnius, Lithuania	Henryk Majchrzak	Katowice, Poland
Ryszarda Chazan	Warsaw, Poland	Ewa Małecka-Tendera	Katowice, Poland
Stanislav Czudek	Ostrava, Czech Republic	Stella Nowicki	Memphis, USA
Jacek Dubiel	Cracow, Poland	Alfred Patyk	Gottingen, Germany
Zbigniew Gasior	Katowice, Poland	Palmira Petrova	Yakutsk, Russia
Mowafaq Muhammad Ghareeb	Baghdad, Iraq	Krystyna Pierzchała	Katowice, Poland
Andrzej Gładysz	Wroclaw, Poland	Waldemar Priebe	Houston, USA
Nataliya Gutorova	Kharkiv, Ukraine	Maria Siemionow	Chicago, USA
Marek Hartleb	Katowice, Poland	Vladyslav Smiiianov	Sumy, Ukraine
Roman Jaeschke	Hamilton, Canada	Tomasz Szczepański	Katowice, Poland
Andrzej Jakubowiak	Chicago, USA	Andrzej Witek	Katowice, Poland
Peter Konturek	Saalfeld, Germany	Zbigniew Wszolek	Jacksonville, USA
Jerzy Korewicki	Warsaw, Poland	Vyacheslav Zhdan	Poltava, Ukraine
Jan Kotarski	Lublin, Poland	Jan Zejda	Katowice, Poland

Distribution and Subscriptions:

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

Graphic design / production:

Grzegorz Sztank

www.red-studio.eu

Publisher:

ALUNA Publishing House

ul. Przesmyckiego 29,

05-510 Konstancin – Jeziorna

www.wydawnictwo-aluna.pl

www.wiadomoscilekarskie.pl

www.wiadlek.pl

CONTENTS

ORIGINAL ARTICLES

- Vladyslav V. Smiiianov, Svitlana M. Piven, Natalia O. Dryga, Olha I. Smiiianova, Inna O. Plakhtiienko, Natalia G. Kuchma, Kateryna I. Zhuk
 MEDICAL AND SOCIAL ANALYSIS OF PATIENTS WHO DIED OF SARS-COV-2 IN THE SPECIALIZED INFECTIOUS DEPARTMENT
 OF THE CITY OF SUMY DURING THE FIRST WAVE OF THE EPIDEMIC 891
- Włodzisław Kuliński, Emilia Gryl
 HIPPOThERAPY IN CEREBRAL PALSY – SURVEY RESEARCH 897
- Hanan Diekan Abbas, Ahmed Hassan Kudhair
 THE EFFECT OF COVID-19 INFECTION ON THE INCIDENCE OF THE RENAL ABNORMALITY FOR THE PATIENT ADMITTED
 IN TO THE COMMUNICABLE DISEASE UNITE AT AL-HAKEEM HOSPITAL 907
- Iryna R. Vyshnevskaya, Tatyana Storozhenko, Mykola P. Kopytsya, Natalia V. Bila, Andrii Kis, Mohammad Kaaki
 THE ROLE OF BIOMARKER MACROPHAGE MIGRATION INHIBITORY FACTOR IN CARDIAC REMODELING PREDICTION IN PATIENTS
 WITH ST-SEGMENT ELEVATION MYOCARDIAL INFARCTION 911
- Vladyslav A. Smiiianov, Leonid B. Markin, Tetiana V. Fartushok, Svitlana R. Slavitych
 EFFECTIVENESS OF THE MAGNESIUM APPLICATION IN OBSTETRICAL PRACTICE 920
- Maksym Herasymiuk, Andrii Sverstiuk, Iryna Kit
 MULTIFACTOR REGRESSION MODEL FOR PREDICTION OF CHRONIC RHINOSINUSITIS RECURRENCE 928
- Anastasiia Efimenko, Oksana Ishchenko, Oleksii Stepanskyi, Dmytro Stepanskyi
 MICROBIOLOGICAL FEATURES OF *STAPHYLOCOCCUS* ASSOCIATED WITH COMPLICATIONS OF DENTAL IMPLANTATION 936
- Basma Adel Ibrahim, Sawsan Ali Hussein, Wasnaa Hadi Abdullah
 COGNITIVE FUNCTIONS IN CHILDREN WITH TYPE I DIABETES 944
- Saad H. Al-Badry, Nowar Gassan Ibrahim, Haider Abdulkareem AlMashhadani, Mustafa M. Kadhim
 THE ROLE OF OXIDATIVE STRESS IN PATIENTS WITH CHRONIC KIDNEY DISEASE 951
- Valeriy Zub, Elina Manzhali
 ASSESSMENT OF QUALITY OF LIFE IN PATIENTS WITH GASTRIC CANCER IN UKRAINE 956
- Svitlana I. Chepyshko, Oleh O. Maksymiv, Mykola L. Gomon, Vitaliy I. Rozhko, Stepan V. Tkachyk, Halyna Yu. Tsybaliuk, Larysa Ya. Fedoniuk
 RESULTS OF INFLAMMATORY AND IMMUNOLOGICAL PARAMETERS OF THE ORAL CAVITY AFTER CYSTECTOMY WITH DIFFERENT
 METHODS OF CONNECTING THE EDGES OF THE OPERATED AREA 964
- Tetiana B. Kodlubovska, Liudmyla V. Piankivska, Volodymyr V. Krasnov, Larisa Yu. Babintseva
 RELATIONSHIP BETWEEN PROFESSIONAL MALADAPTATION OF DOCTORS – INTERNS WITH THE MEANINGFULNESS OF LIFE 972

Ahmed Khalaf Jasim, Dhiaa Jabir Aljuburi, Ali Abbar Mazael, Osamah Aldulhasan, Tamam Alaa Abdulzahra EVALUATION OF VEST-OVER-PANT TECHNIQUE IN THE TREATMENT OF POST-HYPOSPADIAS URETHROCUTANEOUS FISTULA	978
Iryna M. Nikitina, Anna A. Synkina, Yulia B. Yakymchuk, Natalia P. Sukhostavets, Tetiana V. Kopytsia, Svitlana F. Herasymenko, Tetiana V. Babar MENSTRUAL DISORDER IN ADOLESCENTS DURING WAR	984
Viktor M. Shevchuk, Oleg L. Musiienko, Mykyta O. Sokolenko CRIMINAL OFFENCES RELATED TO ILLICIT TRAFFICKING IN FALSIFIED MEDICINES: INVESTIGATION PROBLEMS	992
Mahmood Jawad, Karam Al-Akkam, Manar Mohammed, Saif M. Hassan ROLE OF DIMETHYL FUMARATE (NRF2 ACTIVATOR) IN REDUCING OF CIPROFLOXACIN-INDUCED HEPATOTOXICITY IN RATS VIA THE NRF2/HO-1 PATHWAY	1001
Liudmyla M. Tantsura, Olena Yu. Pylypets, Dmytro V. Tretiakov, Yevhen O. Tantsura VARIANTS OF THE FORMATION AND COURSE OF DRUG-RESISTANT EPILEPSY IN CHILDREN WITH GENETIC POLYMORPHISMS OF <i>CYP2C9</i> , <i>CYP2C19</i> , <i>CYP3A4</i>	1007
Mykhailo S. Myroshnychenko, Nana M. Pasiyeshvili, Nataliia V. Kapustnyk, Volodymyr D. Markovskiy, Yuliia M. Kalashnyk-Vakulenko, Olga V. Zaytseva, Maryna A. Bondarenko, Oleksandr V. Morozov, Dmytro V. Molodan MORPHOLOGICAL ASSESSMENT OF THE LUNGS IN POST-COVID-19 SYNDROME: ANALYSIS OF AUTOPSY MATERIAL	1014
Vitalii V. Ilika, Oleksandra V. Garvasiuk, Oleksandra I. Dogolich, Batih V. Iryna THE FEATURES OF LIMITED PROTEOLYSIS IN PLACENTAL FIBRINOID IN COMBINATION WITH INFLAMMATION AND IRON DEFICIENCY ANEMIA OF PREGNANT WOMEN	1022
Vira Y. Tseluyko, Larysa M. Yakovleva, Olha V. Radchenko, Andriy V. Zhadan, Svitlana M. Sukhova, Oleksandr M. Gurov, Roman N. Askierov PECULIARITIES OF THE COURSE OF PULMONARY EMBOLISM DURING THE COVID-19 PANDEMIC	1029
Zahraa N. Fakhreldain, Hayder Ch. Assad EVALUATION THE PREVALENCE OF MULTIDRUG RESISTANCE BACTERIA AMONG IRAQI PATIENTS AND ITS ASSOCIATION WITH PATIENTS' PREDICTIVE FACTORS: A CROSS-SECTIONAL STUDY	1039
Iryna Khomenko, Oleksandra Ivakhno, Yaroslav Pershehuba, Nataliia O. Serohina TRANSFORMATION CHANGES OF PUBLIC HEALTHCARE WITHIN EPIDEMIOLOGICAL RISK MANAGEMENT LEVEL	1049
Borys Palamar, Svitlana Palamar, Tetiana Miyer, Victoria Zhelanova, Sergey Tukaiev, Hennadii Bondarenko, Natalya Holota STUDYING ANXIETY AS A PREDICTOR IN STUDENTS TO PREDICT THE DEVELOPMENT OF BURNOUT	1054
REVIEW ARTICLES	
Klaudia Ból, Aleksandra Brudzińska, Łukasz Wybrańczyk, Adam Właszczuk EFFECT OF HORMONAL CONTRACEPTION ON DEPRESSION IN WOMEN	1062

Iryna Iakovets, Anatolii Stepanyuk, Valentyna Zverkhovska PENITENTIARY MEDICINE IN THE CONTEXT OF NATIONAL HEALTH CARE REFORM IN UKRAINE	1068
Oleksandr A. Halushko, Olga V. Protsiuk, Oksana I. Pogorila, Maksym M. Synytsyn HYPOGLYCEMIA IN PATIENTS WITH COVID-19: A COINCIDENCE OR A TREND?	1075
Nataliia M. Akhtyrska, Olena Yu. Kostiuchenko STATE REGISTRY OF HUMAN GENOMIC INFORMATION IN UKRAINE: HOW TO ACHIEVE A BALANCE BETWEEN LAW, MEDICINE AND ETHICS	1083
Victoriia V. Haltsova, Sergiy O. Kharytonov, Andrii M. Iashchenko HARM TO LIFE AND HEALTH AS A RESULT OF ACTS OF ILLEGAL ADOPTION: CRIMINAL LEGAL AND MEDICO-PSYCHOLOGICAL ASPECTS	1090
Iryna Iakovets, Olena Haltsova, Oksana Kashyntseva ENSURING THE RIGHT TO HEALTH CARE FOR GROUPS VULNERABLE TO HIV IN PENITENTIARY INSTITUTIONS AND PRE-TRIAL DETENTION CENTERS IN UKRAINE	1097
Larysa Grynko, Hennadii Chorny, Maryna Kostenko SALE OF FALSIFIED MEDICINES VIA THE INTERNET IN UKRAINE: PROBLEMS OF DETECTION AND COUNTERACTION	1106
Halyna V. Bilavych, Natalia V. Sultanova, Iryna S. Litvinenko, Jan V. Biławicz, Nadiya O. Fedchyshyn, Larysa Ya. Fedoniuk, Borys P. Savchuk ANTI-ALCOHOL EDUCATION OF CHILDREN, YOUTH AND ADULTS AS A FACTOR IN THE FORMATION OF HEALTH-SAVING COMPETENCE THROUGH THE PRISM OF THE HISTORICAL EXPERIENCE OF UKRAINE	1113
CASE STUDIES	
Natalya B. Gubergrits, Tetiana L. Mozhyna, Oleksii V. Tsys, Venera S. Rakhmetova GASTRIC AND DUODENAL AMYLOIDOSIS IN AN HIV-INFECTED PATIENT: A CASE REPORT AND A LITERATURE REVIEW	1121
LETTER TO EDITOR	
Julia Maya Nowak A NEW OUTLOOK ON PATIENT CARE – MY EXPERIENCE AT MAYO CLINIC AS AN INTERNATIONAL MEDICAL STUDENT	1130

ORIGINAL ARTICLE

RESULTS OF INFLAMMATORY AND IMMUNOLOGICAL PARAMETERS OF THE ORAL CAVITY AFTER CYSTECTOMY WITH DIFFERENT METHODS OF CONNECTING THE EDGES OF THE OPERATED AREA

DOI: 10.36740/WLek202305112

Svitlana I. Chepyshko¹, Oleh O. Maksymiv¹, Mykola L. Gomon², Vitaliy I. Rozhko¹, Stepan V. Tkachyk¹, Halyna Yu. Tsymbaliuk³, Larysa Ya. Fedoniuk³

¹BUKOVINIAN STATE MEDICAL UNIVERSITY, CHERNIVTSI, UKRAINE

²NATIONAL PIROGOV MEMORIAL MEDICAL UNIVERSITY, VINNYTSIA, UKRAINE

³I. HORBACHEVSKY TERNOPIL NATIONAL MEDICAL UNIVERSITY, TERNOPIL, UKRAINE

ABSTRACT

The aim: To analyze the results of inflammatory and immunological parameters of the oral cavity after cystectomy with different methods of connecting the edges of the operated area.

Materials and methods: The research was conducted in 87 patients who sought surgical treatment of odontogenic cysts of the jaws. Patients were divided into groups depending on the method of wound closure after surgery. We analyzed the results of laboratory tests (leukocytes, ESR, IL-1 β , IL-6, TNF- α , IL-8, NO synthase, MMP-9).

Results: The analysis of the effectiveness of using different methods of approximation of the oral mucosa wound on inflammatory and immunological parameters showed that when using welding of the edges of the operated area using the EKVZ-300 "Patonmed" apparatus, we get positive results of laboratory indicators of inflammatory markers faster (leukocyte counts are normal on day 30, ESR, IL-1 β – on the 14th day, TNF- α – on the 7th day, IL-6, IL-8, NO synthase, MMP-9 – on the 30th day), that is, healing in such patients is faster than in patients with wound closure with classic suture material or laser.

Conclusions: When comparing the methods of approximation of postoperative oral mucosa wounds by different methods, based on the results of these inflammatory and immunological parameters, the best results were obtained when using electric welding of tissues. Further research and use of the proposed method will facilitate and shorten the rehabilitation period of patients after surgery.

KEY WORDS: odontogenic cyst, cystectomy, electric welding of tissues, laboratory tests

Wiad Lek. 2023;76(5 p.1):964-971

INTRODUCTION

Odontogenic cysts of the jaws are one of the most common diagnoses requiring surgical treatment. Among them, radicular cysts are the most common. They are the result of an inflammatory process in the periapical tissues associated with necrotic and infected pulp [1]. Humoral and cellular immune responses play a central role in the pathogenesis of these lesions. The most important role in the growth of these lesions is played by the proinflammatory cytokines TNF-alpha, IL-1, and IL-6 [2]. The cytokine can be secreted by macrophages, monocytes, and other cells of the immune system and may be involved in skeletal homeostasis, including osteoclast formation, and bone resorption in the maxillofacial region [3]. For a more in-depth study of this issue, it is necessary to first give a generalized overview of the mechanism of growth

of maxillofacial radicular cysts with an indication of the clinical aspect, as well as to clarify the role of cytokines in this pathophysiological process [4, 5].

Radicular cysts are the most common among jaw cysts, and, according to various authors [6, 7, 8], they account for 6 to 84 % of all periapical lesions. At the same time, tooth-containing cysts (follicular cysts) and keratocysts are diagnosed in 37 % and 6-32 % of patients, respectively. When studying the prevalence of radicular cysts by gender, a small difference was found between men and women. According to some authors, odontogenic cysts were more commonly detected in men (55.9-65.0 %) than in women (44.1-35.0 %) [9, 10]. Other authors report that odontogenic cysts are more common in women (52.37-65.0 %) and less common in men (35.0-47.62 %) [7].

Surgical operations performed at present and associated with cystectomy are based on two main points: tissue separation and subsequent connection [11, 12]. One of the conditions for a properly performed operation is the ability to achieve rapid wound healing, provide a good cosmetic effect, and prevent complications [13, 14, 15]. During many surgical interventions in the oral cavity, the failure of a single suture in the postoperative period can lead to significant complications that usually require repeated operations [16, 17, 18, 19].

THE AIM

To analyze the results of inflammatory and immunological parameters of the oral cavity after cystectomy with different methods of connecting the edges of the operated area.

MATERIALS AND METHODS

In accordance with the aim, laboratory tests were performed in patients after surgical treatment of odontogenic cysts of the jaws at the Educational-Treatment Center "University Clinic" (Chernivtsi, Ukraine).

The patients were divided into three groups: Group I (20 patients) – cystectomy with wound suturing using suture material «Vicryl»; Group II (28 patients) – cystectomy with wound edge consolidation using the laser scalpel LSP – "IRE-POLUS" and Group III (39 patients) – the edges of the postoperative defect were connected by using the electric welding apparatus EKVZ-300 "Patonmed".

After cystectomy, 20 patients (group A, 1 control) had their surgical wounds sutured with suture material «Vicryl». In 28 patients (group B, 2 control), after cystectomy, the wound edges were approximated using the LSP laser – "IRE-POLUS", with an irradiation wavelength of $0.97 \pm 1.56 \mu\text{m}$, maximum irradiation power of 30/60 W; continuous, pulsed, pulsed-periodic mode of operation [20]. A laser with a wavelength of $1.06 \mu\text{m}$, a power of 0.71-0.85 W, a continuous irradiation mode, a light guide diameter of 0.6 mm was used to join the wound edges in the area of contact by moving the source along the surface of the wound edge at a distance of 1.0-3.0 mm, with a light guide movement speed of no more than 1.0 mm/sec [21].

In 39 patients (group C, main), the edges of the postoperative wound, after cystectomy, were welded using the apparatus EKVZ-300 "Patonmed".

Patients in this group were treated with the wound edges using the apparatus EKVZ-300 "Patonmed" in the automatic welding mode N2 using the following parameters: voltage – 25 W, duration of exposure to

high-frequency current on tissues no more than 2-6 s/ per point, duration of voltage amplification – 150 ms, specific pressure force on tissues in the range of 0.3-0.7 N/cm², current frequency – 66 kHz, temperature in the exposure zone 55-65 °C. Patients of all research groups were given recommendations for wound care in the postoperative period and prescribed drug therapy: nimesulide – 100 mg, 1 sachet twice a day for three days after the intervention; rinsing the oral cavity with chlorhexidine 0.06 % after each meal; topical cold for 5 minutes three times a day for the first three days after the intervention.

RESULTS

As a result of our studies, we found that in healthy subjects of the control group, the content of leukocytes in the oral fluid was $276.80 \pm 45.72 \times 10^9/\text{L}$ (Table I). On the first postoperative day, the content of leukocytes in the oral fluid increased in patients of all Patients of all research groups compared with the data in the control group: by 63.87 %, $p < 0.05$ in group A, by 100 %, $p < 0.01$ in group B, and by 81.94 % in group C. At the same time, the values of the obtained indicators did not differ in statistical significance in the intergroup comparison. On the 5th postoperative day, the quantitative content of leukocytes in the oral fluid in the patients of all research groups continued to increase and was significantly higher than in the control group in all Patients of all research groups, $p < 0.01$. However, the minimum values were determined in patients of groups C and A – $523.90 \pm 61.80 \times 10^9/\text{L}$ and $554.20 \pm 60.15 \times 10^9/\text{L}$, respectively, $p < 0.01$, $p_1 > 0.05$. The maximum value was studied in patients of group B – $763.90 \pm 62.18 \times 10^9/\text{L}$, which was, on average, 29.43 % higher than in patients of groups A and C, $p_1 < 0.05$, $p_2 < 0.05$.

On day 7th, 14th and 30th day of postoperative follow-up, a moderate decrease in the number of leukocytes in the oral fluid was determined in patients of the research groups, which, however, remained significantly higher than in patients of control group B, $p < 0.01$, $p < 0.05$, than in patients of groups A and C, $p_1 > 0.05$, $p_2 < 0.05$, on average.

The investigation of ESR parameters in the research groups (Table II) revealed a tendency to increase the data of this parameter on the 1st-5th postoperative day, which reached maximum values on the 5th day of observation and was higher than in the control group: in 3.3 times in patients of group A; in 3.7 times in patients of group B and 3.0 times in patients of group C, $p < 0.01$; p_1 , $p_2 > 0.05$. It should be noted that on the 7th day of postoperative follow-up in patients of group C, the ESR was minimal ($8.34 \pm 1.58 \text{ ml/h}$) and equal to that of the control group, p , p_1 , $p_2 > 0.05$. At the same time, in

Table I. Dynamics of the number of leukocytes in the oral fluid in the patients of the patients of all research groups at different follow-up periods

Research groups	1st day	5th day	7th day	14th day	30th day
Control group	276.80±45.72 x 10 ⁹ /l				
Group A	453.60± ±56.20 [°]	554.20± ±60.15 [°]	454.0± ±56.20 ^{°°}	353.50± ±51.18	273.50± ±49.82
Group B	553.60± ±58.42 [°]	763.90± ±62.18 ^{°,**}	603.70± ±57.83 [°]	503.20± ±50.00 ^{**}	400.70± ±49.97
Group C	503.60± ±57.28 [°]	523.90± ±61.80 ^{°,ΔΔ}	423.70± ±55.19 ^{°°,ΔΔ}	323.00± ±49.86ΔΔ	272.50± ±48.22

Notes:

1. [°]p<0.01; ^{°°}p<0.05 – significant difference in values relative to the average statistical norm.2. ^{**}p1<0.05 – significant difference in values relative to group A.3. ^{ΔΔ}p2<0.05 – significant difference in values relative to group B.**Table II.** Changes in ESR in patients of the research groups in different postoperative periods

Research groups	1st day	5th day	7th day	14th day	30th day
Average statistical norm	5.5±1.10 mm/h				
Group A	13.28±1.46 [°]	18.33±1.72 [°]	10.56±1.62 ^{°°}	7.23±1.60	5.00±1.21
Group B	14.39±1.49 [°]	20.45±1.98 [°]	12.55±1.70 [°]	9.20±1.66	7.18±1.24
Group C	13,84±1.47 [°]	16.42±1.83 [°]	8.34±1.58	5.29±1.57	5,10±0.93Δ

Notes:

1. [°]p<0.01; ^{°°}p<0.05 – significant difference in values relative to the data in the control group.2. ^Δp2<0.05 – significant difference in values relative to the data in group B.**Table III.** Dynamics of IL-1β content in the oral fluid of patients after cystectomy at different follow-up periods

Research groups	1st day	5th day	7th day	14th day	30th day
Control group	68.34±6.03 pg/ml				
Group A	98.65± ±6.68 [°]	118.76± ±9.13 [°]	88.46± ±6.13 ^{°°}	78.63± ±6.12	68.40± ±6.11
Group B	110.20± ±6.90 [°]	130.42± ±6.15 [°]	100.12± ±6.14 ^{°,**}	90.00± ±6.13 ^{°°}	75.15± ±6.13
Group C	104.42± ±6.69 [°]	114.62± ±6.14 [°]	84.32± ±6.12ΔΔ	64.12± ±6.11ΔΔ	64.50± ±6.12

Notes:

1. [°]p<0.01; ^{°°}p<0.05 – significant difference in values relative to the control group.2. ^{**}p1<0.05 – significant difference in values relative to the data of group A.3. ^{ΔΔ}p2<0.05 – significant difference in values relative to the data of group B.

patients of groups A and B, the value was in 1.9 times, p<0.05, and in 2.3 times, p<0.01 higher, respectively, compared to the control data. On the 7th-14th day of observation, the ESR value decreased compared with the control data, p>0.05, with the most pronounced positive dynamics in patients of group C.

The early action of interleukin IL-1β plays a key role in the development and regulation of innate and acquired immunity, being one of the first to be involved in the defense response to pathogenic factors, activating neutrophils, and stimulating the synthesis of acute phase proteins [12, 22].

It was established (Table III) that on the 1st postoperative day, the content of IL-1β in the oral fluid of

patients of the research groups increased compared to the data in the control group: by 44.35 % in group A, by 61.25 % in group B, and by 52.79 % in group C, p<0.01, p1, p2>0.05.

On the 5th postoperative day, the maximum increase in values was noted relative to the data in the control group: by 73.78 % in group A, by 90.83 % in group B and by 67.72 % in group C, p<0.01, p1, p2>0.05. It should be noted that on the 7th, 14th, 30th postoperative days, the level of IL-1β in the oral fluid of the subjects decreased to the maximum and reached normal levels in the patients of group C.

The concentration of the proinflammatory cytokine IL-6, which plays a leading role in the induction of acute

phase proteins [24], increased in the oral fluid of the subjects on the 1st day after cystectomy compared with the control data: in 3.0 times in group A, in 3.3 times in group B, $p_1 < 0.05$, and in 3.1 times, $p_1, p_2 > 0.05$, (Table IV).

On the 5th day of the postoperative period, in groups A and B, the level of IL-6 in the oral fluid increased and was higher than in the control group by 3.5 times and 3.8 times, $p_1 < 0.05$, respectively, $p < 0.01$. At the same time, in patients of group C, a decrease in the content of IL-6 in the oral fluid was studied relative to the data

of the previous research period, but the obtained values remained in 2.7 times higher than in the control group, $p < 0.01$. It should be noted that at this time of the investigation, the content of IL-6 in the oral fluid of patients in group C was, on average, in 1.3 times lower than in patients in groups A and B. On the 7th, 14th and 30th days of observation in patients of group C, IL-6 in the oral fluid did not differ statistically from the data in the control group, $p > 0.05$, and was lower in patients of groups A and B.

Table IV. Dynamics of IL-6 content in the oral fluid of patients after cystectomy in different periods of observation

Research groups	1st day	5th day	7th day	14th day	30th day
Control group	2.20±0.10 pg/ml				
Group A	6.65± ±0.19°	7.76± ±0.21°	6.00± ±0.15°	3.48± ±0.12°	2.95± ±0.10°
Group B	7.29± ±0.21°,**	8.40± ±0.23°,**	6.49± ±0.16°,**	4.00± ±0.13°,**	3.15± ±0.11°
Group C	6.90± ±0.20°	6.00± ±0.17°*,Δ	4.82± ±0.13°*,Δ	2.80± ±0.10°*,Δ	2.10± ±0.09*,Δ

Notes:

1. ° $p < 0.01$ – significant difference in values compared to the control group.
2. * $p_1 < 0.01$; ** $p_1 < 0.05$ – significant difference in values relative to the data of group A.
3. Δ $p_2 < 0.01$ – significant difference in values relative to the data of group B.

Table V. Dynamics of TNF-α content in the oral fluid of patients after cystectomy at different follow-up periods

Research groups	1st day	5th day	7th day	14th day	30th day
Control group	89.70±13.40 pg/ml				
Group A	109.90± ±14.20	180.00± ±15.43°	166.55± ±14.82°	100.36± ±14.10	93.26± ±23.10
Group B	99.80± ±12.65	160.63± ±14.80°	140.80± ±14.20°°	92.82± ±13.25	89.75± ±22.80
Group C	104.85± ±14.17	144.95± ±14.47°°	90.22± ±14.00*,ΔΔ	89.50± ±12.48	89.50± ±22.76

Notes:

1. ° $p < 0.01$; ° $p < 0.05$ – significant difference in values compared to the control group.
2. * $p_1 < 0.01$ – significant difference in values compared to the data in group A.
3. Δ $p_2 < 0.05$ – significant difference in values relative to the data in group B.

Table VI. Dynamics of IL-8 content in the oral fluid of patients after cystectomy in different periods of observation

Research groups	1st day	5th day	7th day	14th day	30th day
Control group	5.77±0.67 pg/ml				
Group A	16.28± ±0.82°	14.00± ±0.75°	13.80± ±0.73°	10.50± ±0.48°	8.25± ±0.30°
Group B	15.24± ±0.77°	13.14± ±0.73°	12.26± ±0.69°	9.15± ±0.46°,**	8.36± ±0.31°
Group C	16.00± ±0.80°	13.80± ±0.71°	9.45± ±0.54°,*	6.28± ±0.31*,Δ	5.72± ±0.29*,Δ

Notes:

1. ° $p < 0.01$ – significant difference in values compared to the control group.
2. * $p_1 < 0.01$; ** $p_1 < 0.05$ – significant difference in values relative to the data of group A.
3. Δ $p_2 < 0.01$ – significant difference in values relative to the data of group B.

Table VII. Dynamics of NO synthase in the oral fluid of patients of the research groups at different observation periods

Observation periods	Research groups	NO ₂ , μmol/l	NO ₃ , μmol/l	NO ₂ + NO ₃ , μmol/l
1 day	Control group	13.08±2.18	22.50±3.25	35.58±3.81
	Group A	10.15±1.26	17.46±2.18	27.61±3.44
	Group B	9.24±1.15	16.35±2.00	25.59±3.15 ^{oo}
	Group C	9.70±1.18	16.91±2.15	26.61±3.33
5 day	Group A	7.95±1.14 ^{oo}	8.73±1.36 ^o	16.68±2.50 ^o
	Group B	5.94±0.92 ^o	8.18±1.30 ^o	14.12±2.22 ^o
	Group C	6.95±0.98 ^{oo}	11.27±1.82 ^o	18.22±2.80 ^o
7 day	Group A	9.05±1.10	9.84±1.50 ^o	18.89±2.60 ^o
	Group B	7.14±0.99 ^{oo}	9.29±1.48 ^o	16.43±2.47 ^o
	Group C	9.15±1.10	13.50±1.95 ^{oo}	22.65±3.05 ^{oo}
14 day	Group A	10.20±1.24	14.89±2.00 ^{oo}	25.09±3.24 ^{oo}
	Group B	9.36±1.12	15.34±2.13	24.70±3.25 ^{oo}
	Group C	12.48±1.20	18.03±2.62	30.51±3.82
30 day	Group A	12.00±1.21	18.92±2.66	30.92±3.87
	Group B	11.58±1.19	17.67±2.47	29.25±3.66
	Group C	13.62±1.22	21.36±2.92	34.98±4.14

Notes. ^op<0.01; ^{oo}p<0.05 – significant difference in values relative to the control data.

Table VIII. Dynamics of MMP-9 content in the oral fluid of patients of the research groups at different times of the postoperative period

Research groups	Observation periods				
	1st day	5th day	7th day	14th day	30th day
Group A	4.780± ±0.956 ^o	10.523± ±2.104 ^o	3.507± ±0.701 ^o	2.406± ±0.481 ^o	1.394± ±0.278 ^{oo}
Group B	6.200± ±1.240 ^o	11.634± ±2.326 ^o	5.817± ±1.163 ^o	4.706± ±0.941 ^{o,ΔΔ}	2.582± ±0.516 ^{o,ΔΔ}
Group C	3.990± ±0.798 ^o	9.745± ±1.949 ^o	2.784± ±0.556 ^{o,**}	1.673± ±0.334 ^{oo,*}	0.773± ±0.154 [*]
Control group	0.764±0.130 pg/ml				

Notes:

1. ^op<0.01; ^{oo}p<0.05 – significant difference in values relative to the control group.

2. ΔΔp1<0.05 – significant difference in values relative to the data of group A.

3. *p2<0.01; **p2<0.05 – significant difference in values compared to the data of group B.

In practically healthy subjects of the control group, the content of TNF-α in the oral fluid was at a low level (89.70±13.40 ng/ml), which probably provides only the physiological value of direct and feedback signals in the cytokine system, maintaining normal regulation of immunoreactivity (Table V). The increase in this indicator occurs under the influence of antigens of viruses or bacteria in response to the inflammatory process [22, 23].

On the 1st postoperative day after cystectomy, an increase in the level of TNF-α in the oral fluid was observed in patients of the research groups, on the 5th day of observation – a significant increase in this parameter in the research groups relative to the values in the control: by 100.66 % in group A, by 79.07 % in group B, p<0.01, p1>0.05; and by 61.59 % in group C, p<0.05, p1, p2>0.05.

On the 7th day, in patients of group C, the concentration of TNF-α in the oral fluid did not differ statistically from that in the control group, p>0.05, and was lower than in patients of groups A and B by 84.60 % and 56.06 %, respectively.

The content of IL-8, which is produced under the influence of bacterial endotoxins and cytokines, mainly TNF-α and IL-1 [24, 25], increased in the oral fluid of the subjects on the 1st postoperative day compared to the control data: in 2.8 times in patients of group A, in 2.6 times in patients of group B, and 2.7 times in patients of group C, p<0.01; p1, p2>0.05. On the 5th day of the research, a gradual decrease in the content of IL-8 was observed, on the 7th day of postoperative observation, the minimum values were studied in patients of group C – 9.45±0.54 ng/ml, p, p1, p2<0.01, which were 1.5 times and 1.3 times, p1>0.05, lower compared with those

in patients of groups A and B, respectively, $p < 0.01$. On the 14th day of the postoperative research in patients of group C, the concentration of IL-8 in the oral fluid did not differ significantly from that in the control group, $p > 0.05$ (Table VI).

On the 1st day of postoperative intervention in patients of the research groups, after cystectomy, the decrease in the content of NO synthase in the oral fluid was studied in relation to the data in the control group. At the same time, the maximum decrease in these parameters was determined in patients of groups B and C, on average, by the level of NO_2 – by 22.60 %, NO_3 – by 26.10 %, and $\text{NO}_2 + \text{NO}_3$ – by 26.64 %, $p > 0.05$, $p_1 > 0.05$. On the 5th postoperative day, a further decrease in the studied levels of indicators in the oral fluid was determined. On the 7th day of the postoperative period, positive dynamics of NO_2 content in the oral fluid of subjects of groups A and C was noted, relative to the data in the control group, $p > 0.05$ (Table VII).

After 2 weeks of observation (14 days) in patients of group C, the data on the content of NO synthase in the oral fluid were normalized and equal to the values in the control group, $p > 0.05$.

It should be added that during the study at all periods of observation, there was no significant difference in the values of NO synthase in the oral fluid in the intergroup comparison, $p_1, p_2 > 0.05$.

It was found that in the control group, the content of MMP-9 in the oral fluid (Table VIII) was 0.764 ± 0.130 pg/ml. At the same time, in patients of the research groups on the 1st postoperative day, an increase in the values of the parameter was established relative to the data in the control group. On the 5th day of postoperative observation, the maximum increase in the level of MMP-9 in the oral fluid was studied.

At the same time, on the 7th-14th days of the investigation, the content of MMP-9 in the oral fluid decreased, but remained significantly higher than in the control group after two weeks of observation. At the end of the research (the 30th day), in patients of group C, who underwent electric welding of the edges of the postoperative wound, the content of MMP-9 in the oral fluid was equal to that of the control group, $p > 0.05$, $p_2 < 0.01$.

DISCUSSION

One of the most common interventions in the clinic of surgical dentistry, which ranks second after tooth extraction and accounts for more than 40 %, is the removal of odontogenic cysts [26]. The share of radicular cysts among the total number of maxillofacial cysts accounts for 52 % [27]. Modern advances in medicine allow the use of fundamentally new methods of treatment of odontogenic cysts in order to improve the results of treatment and subsequent rehabilitation [28, 29]. The method we propose is based on the use of tissue electric welding technology to approximate the wound edges after cystectomy.

Analysis of the effectiveness of different methods of approximation of the oral mucosa wound on inflammatory and immunological parameters showed that on the first postoperative day the content of leukocytes in the oral fluid increased in patients of all research groups compared to the data in the control group, $p < 0.05$; 0.01 . At the same time, the values of the obtained indicators did not differ in statistical significance in the intergroup comparison. On the 5th postoperative day, the quantitative content of leukocytes in the oral fluid continued to increase and was significantly higher than in the control group in all research groups, $p < 0.01$. However, the minimum values of the analyzed parameter were determined in the patients of groups C and A – $523.90 \pm 61.80 \times 10^9/\text{L}$ and $554.20 \pm 60.15 \times 10^9/\text{L}$, respectively, $p < 0.01$, $p_1 > 0.05$. The maximum values of this parameter were studied in the patients of group B – $763.90 \pm 62.18 \times 10^9/\text{L}$, which was, on average, by 29.43 % higher than in the patients of groups A and C, $p_1 < 0.05$, $p_2 < 0.05$. In 1 month after cystectomy in the patients of group A, where the edges of the postoperative wound were sutured with «Vicryl», and in patients of group C, when using the device ECVZ-300 «Patonmed», the quantitative content of leukocytes in the oral fluid was equal to that in the control group, $p > 0.05$. At the same time, when approximating the edges of the oral mucosa with a laser, in patients of group B, the leukocyte content was by 36.26 % higher than in patients of the other groups, on average, $p, p_1, p_2 > 0.05$. Similar data were obtained by analyzing other indicators.

In a previous research [30] related to this problem, we studied the dynamics of clinical parameters when using different methods of approximating the edges of the oral mucosa. The results showed the effectiveness of the proposed method using electric welding of tissues.

CONCLUSION

When comparing the methods of approximation of postoperative oral mucosa wounds by different methods, based on the results of these inflammatory and immunological parameters, the best results were obtained using electric welding of tissues. In the surgical treatment of odontogenic cysts by the proposed method, a decrease in inflammatory and immunological reactions in the oral fluid of the subjects was found, which was confirmed by a decrease in the levels of IL-1 β , $p - p_1 > 0.05$, $p_2 < 0.05$, TNF – L, $p > 0.05$, $p_1 - p_2 < 0.01$, on the 7th day of observation; quantitative content of leukocytes, $p - p_1 > 0.05$, $p_2 < 0.05$, levels of IL – 8, $p > 0.05$, $p_1 - p_2 < 0.01$, activity of NO – synthase, $p - p_2 > 0.05$ on the 14th day of research; IL concentrations – 6, $p > 0.05$, $p_1 - p_2 < 0.01$, MMP activity – 9, $p - p_2 > 0.05$ on the 30th day after treatment and ESR parameters on the 7th day of observation, $p - p_2 > 0.05$.

REFERENCES

1. Bilodeau EA, Collins BM. Odontogenic Cysts and Neoplasms. *Surg Pathol Clin*. 2017;10(1):177-222. doi: 10.1016/j.path.2016.10.006
2. Neto ND, Porpino MTM., Antunes HD, Rodrigues RCV, Perez AR, Pires FR, et al. Pro-inflammatory and anti-inflammatory cytokine expression in post-treatment apical periodontitis. *Journal of Applied Oral Science*. 2018; 4:55. doi: 10.1590/1678-7757-2017-0455
3. Chen JH, Tseng CH, Wang WC, Chen CY, Chuang FH, Chen YK. Clinicopathological analysis of 232 radicular cysts of the jawbone in a population of southern Taiwanese patients. *Kaohsiung J Med Sci*. 2018;34(4):249-54. doi: 10.1016/j.kjms.2018.01.011
4. Mokryk O.YA., Havryl'tsiv S.T., Korniyenko M.M., Petrov B.P. Suchasni khirurhichni metody likuvannya odontohennykh kist shchelep [Modern surgical methods of treatment of odontogenic cysts of the jaws]. *Ukrainian Journal of Medicine, Biology and Sports*. 2021;6(6):8-19. doi: 10.26693/jmbs06.06.008 (in Ukrainian)
5. Pantus A, Rozhko M, Yarmoshuk I, Kohut V, Solodzhuk YU. Analiz khirurhichnykh metodyk ta pidkhodiv usunennya defektiv kistok lytsevoho skeletu [Analysis of surgical techniques and approaches to eliminate bone defects of the facial skeleton]. *Journal «Art of Medicine»*. 2019; 1:100-104. (in Ukrainian)
6. Havryl'tsiv S.T. (2020) Vdoskonalennya diahnostychno-likuval'noho alhorytmu pry radykulyarnykh kistakh shchelep patsiyentiv iz dyferentsiyovanyim osteoreheneratorynym statusom [Improvement of the diagnostic and therapeutic algorithm in radicular cysts of the jaws of patients with differentiated osteoregenerative status] Extended abstract of candidate's thesis. Lviv. (In Ukrainian).
7. Havryl'tsiv S.T. Suchasni metody likuvannya kistkovykh defektiv shchelep, shcho utvorylysi' pislya vydalennya radykulyarnykh kist [Modern methods of treatment of bone defects of the jaws formed after removal of radicular cysts]. *Journal of Scientific Research*. 2018; 3:5-12. doi: 10.11603/2415-8798.2018.3.9403
8. Malanchuk VO., Shvydchenko V.S., Halatenko N.A., Kulyesh D.V. (2018) Biodehraduyuchi kompozyty dlya usunennya peryradykulyarnykh defektiv kistky ta poperedzhennya zmishchennya zubiv [Biodegradable composites for the elimination of periradicular bone defects and prevention of tooth displacement]. *Materialy Mizhnarodnoyi naukovo-praktychnoyi konferentsiyi "Suchasna stomatolohiya ta shchelepno-lytseva khirurhiya" – Materials of the International Scientific and Practical Conference «Modern Dentistry and Oral and Maxillofacial Surgery»*. (pp 101-104). Kyiv. (In Ukrainian)
9. Tanaka OM, Meira TM, Batista Rodrigues AC, Willems G, Baggio GL, Couto Souza PH. Marsupialization of a Large Radicular Cyst with Extensive Maxillary Tooth Displacement: Eight-Year Follow-Up. *J Dent Child (Chic)*. 2019;86(1):64-8.
10. Tamiolakis P, Thermos G, Tosios KI, Sklavounou-Andrikopoulou A. Demographic and Clinical Characteristics of 5294 Jaw Cysts: A Retrospective Study of 38 Years. *Head Neck Pathol*. 2019;13(4):587-96. doi: 10.1007/s12105-019-01011-7
11. Dragovic M, Pejovic M, Stepic J, Colic S, Dozic B, Dragovic S, et al. Comparison of four different suture materials in respect to oral wound healing, microbial colonization, tissue reaction and clinical features-randomized clinical study. *Clin Oral Investig*. 2020;24(4):1527-41. doi: 10.1007/s00784-019-03034-4
12. Suthar P, Shah S, Waknis P, Limaye G, Saha A, Sathe P. Comparing intra-oral wound healing after alveoloplasty using silk sutures and n-butyl-2-cyanoacrylate. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*. 2020;46(1):28-35. doi: 10.5125/jkaoms.2020.46.1.28
13. Bourdel N, Chauvet P, Roman H, Pereira B, Somcutian O, Dechelotte PJ, et al. Comparison between resection, bipolar coagulation and Plasmajet®: A preliminary animal study. *Eur J Obstet Gynecol Reprod Biol*. 2017; 211:127-33. doi: 10.1016/j.ejogrb.2017.02.004
14. Doroshenko M, Guerra A, Vu L. Airway For Laser Surgery [Internet]. Treasure Island (FL): StatPearls Publishing; 2021 [cited 2022 Oct 18]. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK564406/>
15. Nahaychuk VV, Shkil'nyak LI. Osoblyvosti diahnostyky, likuvannya i profilaktyky patolohichnykh rubtsiv shchelepno-lytsevoyi dilyanky [Features of diagnosis, treatment and prevention of pathological scars of the maxillofacial area]. *Ukrainian dental almanac*. 2017; 2:74-78. (In Ukrainian)
16. Pylypiv N.V., Korniyenko M.M. (2021). Osoblyvosti vyboru khirurhichnoho metodu likuvannya khronichnykh periapikal'nykh zapal'no-destruktyvnykh protsesiv [Features of the choice of surgical treatment of chronic periapical inflammatory and destructive processes]. *Proceedings of the International Scientific and Practical Conference Medical and Pharmaceutical Sciences: Analysis of the Present and Forecast of the Future*. (pp 22-25). Dnipro. (In Ukrainian)
17. Korniyenko M.M. (2016) Udoskonalennya khirurhichnykh metodiv likuvannya khvorykh z khronichnymy periodontytamy ta odontohennymy kistamy shchelep [Improvement of surgical methods of treatment of patients with chronic periodontitis and odontogenic cysts of the jaws]. Extended abstract of candidate's thesis. Lviv. (In Ukrainian).
18. Hong B, Winkel A, Ertl P, Stumpp SN, Schwabe K, Stiesch M, et al. Bacterial colonisation of suture material after routine neurosurgical procedures: relevance for wound infection. *Acta Neurochir (Wien)*. 2018;160(3):497-503. doi: 10.1007/s00701-017-3404-9
19. Li W, Xiao M, Chen Y, Yang J, Sun D, Suo J, et al. Serious postoperative complications induced by medical glue: three case reports. *BMC Gastroenterology*. 2019;19(1):224. doi: 10.1186/s12876-019-1142-6.
20. Gomathi P, Kavitha J, Rajasekar S, Srinivasan S, Lakshmisree S. Treatment of Chronic Inflammatory Gingival Enlargement Using Electrosurgery: A Case Report. *Journal of Dental and Medical Sciences*. 2020;19(3):17-20.
21. Fu W, Wo C. The use of laser in dentistry: a narrative review. *J Biol Regul Homeost Agents*. 2021;35(1 Suppl 1):11-8.
22. Malik A, Chawla S, Tauseef A, Sohail H, Ijaz F, Malik A, et al. Association of Oxidative Stress and Production of Inflammatory Mediators Matrix Metalloproteinase-9 and Interleukin 6: Systemic Events in Radicular Cysts. *Cureus [Internet]*. 2020 [cited 2022 Nov 15];12(4): e7822. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7249770/pdf/cureus-00120000007822.pdf> doi: 10.7759/cureus.7822

23. Alvares PR, de Arruda JAA, Silva LVO, da Silva LP, do Nascimento GJF, da Silveira MMF, et al. Immunohistochemical Analysis of Cyclooxygenase-2 and Tumor Necrosis Factor Alpha in Periapical Lesions. *J Endod.* 2018;44(12):1783-7. doi: 10.1016/j.joen.2018.09.002i
24. Brito LNS, de Lemos Almeida MMR, de Souza LB, Alves PM, Nonaka CFW, Godoy GP. Immunohistochemical Analysis of Galectins-1, -3, and -7 in Periapical Granulomas, Radicular Cysts, and Residual Radicular Cysts. *J Endod.* 2018;44(5):728-33. doi: 10.1016/j.joen.2018.01.008
25. Mangashetti LS, Khapli SM, Wani MR. IL-4 inhibits bone-resorbing activity of mature osteoclasts by affecting NF-kappa B and Ca2+ signaling. *Immunologia.* 2017; 175:917-25. doi: 10.4049/jimmunol.175.2.917
26. Kilinc A, Gundogdu B, Saruhan N, Yalcin E, Ertas U, Urvasizoglu G. Odontogenic and nonodontogenic cysts: An analysis of 526 cases in Turkey. *Niger J Clin Pract.* 2017;20(7):879-83. doi: 10.4103/1119-3077.212448
27. Oliveros-Lopez L, Fernandez-Olavarria A, Torres-Lagares D, Serrera-Figallo MA, Castillo-Oyague R, Segura-Egea JJ, et al. Reduction rate by decompression as a treatment of odontogenic cysts. *Med Oral Patol Oral Cir Bucal [Internet].* 2017[cited 2022 Sep 19];22(5): e643-50. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5694189/pdf/medoral-22-e643.pdf> doi: 10.4317/medoral.21916
28. Kammer PV, Mello FW, Rivero ERC. Comparative analysis between developmental and inflammatory odontogenic cysts: retrospective study and literature review. *Oral Maxillofac Surg.* 2020;24(1):73-84. doi: 10.1007/s10006-019-00816-8
29. Kirsanova NM, Lomakin MV, Soloshchanskii II. General and specific cases of hemostasis in reconstructive surgery of oral cavity. *Journal of Dentistry.* 2018;22(4):212-6. doi: 10.18821/1728-2802-2018-22-4-212-216
30. Chepyshko S.I., Maksymiv O.O. Dynamika klinichnykh pokaznykiv pislya provedennya tsystektomiyi z vykorystannyam elektrozv'yuvannya krayiv operovanoi dilyanky u patsiyentiv z odontohennymy kistamy shchelep. [Dynamics of clinical parameters after cystectomy using electric welding of the edges of the operated area in patients with odontogenic cysts of the jaws]. *Journal «Modern dentistry».* Kyiv, 2022; 1-2:52-56. (In Ukrainian).

The study was conducted as a fragment of a comprehensive research work of the Department of Orthopedic Dentistry of Bukovinian State Medical University «Etiopathogenetic aspects of rehabilitation of major dental diseases of the maxillofacial area», (state registration number 0121U109997.)

ORCID and contribution:

Svitlana Chepyshko: 0000-0001-7008-0622 ^{A, B, C, D}

Oleh Maksymiv: 0000-0003-4975-7031 ^{A, B, C, D}

Mykola Gomon: 0000-0002-3557-2928 ^{E, F}

Vitaliy Rozhko: 0000-0002-0757-9671 ^{B, E, F}

Stepan Tkachyk: 0000-0002-2797-4994 ^{E, F}

Halyna Tsybaliuk: 0000-0003-1452-9442 ^{E, F}

Larysa Fedoniuk: 0000-0003-4910-6888 ^{B, D, F}

Conflict of interest:

The Authors declare no conflict of interest

CORRESPONDING AUTHOR

Larysa Fedoniuk

Medical Biology Department

I. Horbachevsky Ternopil National Medical University

Valova street, 9, Ternopil, 46000, Ukraine

tel: +38(067)3999143

e-mail: Fedonyuk22Larisa@gmail.com

Received: 11.08.2022

Accepted: 20.04.2023

A – Work concept and design, B – Data collection and analysis, C – Responsibility for statistical analysis, D – Writing the article, E – Critical review, F – Final approval of the article

 Article published on-line and available in open access are published under Creative Common Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0)